



Sutab Prep

- Do not take other laxatives while taking Sutab.
- Do not take oral medications within 1 hour of starting each dose of Sutab.
- If taking tetracycline or fluoroquinolone antibiotics, iron, digoxin chlorpromazine, or penicillin, take these medications at least 2 hours before and not less than 6 hours after administration of each dose of Sutab.

The Dosing Regimen

Sutab is a split-dose (2 day) regimen. A total of 24 tablets is required for complete preparation for colonoscopy. You will take the tablets in two doses of 12 tablets each. Water must be consumed with each dose of SUTAB, and additional water must be consumed after each dose.

DOSE 1 - Day Prior to Colonoscopy

Take the tablets with water.

- Open 1 bottle of 12 tablets.
- Fill the provided container with 16 ounces of water (up to the fill line). Swallow each tablet with a sip of water, and drink the entire amount of water over 15 to 20 minutes.
- One hour after the last tablet is taken, fill the container with an additional 16 ounces of water and drink it over the next 30 minutes.
- 30 minutes after finishing the second container of water, drink an additional 16 ounces of water.

It is VERY important to continue drinking water throughout this process.

DOSE 2 - Day of the Colonoscopy

- Continue to consume only clear liquids until 3 hours before scheduled colonoscopy.
- The morning of the colonoscopy (5-6 hours before the scheduled colonoscopy), open the second bottle of 12 tablets.
- Repeat the above steps from DOSE 1

IMPORTANT: You must complete all SUTAB tablets and required water at least 3 hours before colonoscopy.

If you experience symptoms, such as nausea, bloating or cramping, pause or slow the rate of drinking the water until the symptoms diminish.